

NOTE

W.W. Toller · R. Rowan · N. Knowlton

Genetic evidence for a protozoan (phylum Apicomplexa) associated with corals of the *Montastraea annularis* species complex

Received: 8 June 2001 / Accepted: 13 March 2002 / Published online: 8 June 2002
© Springer-Verlag 2002

Keywords Coccidia · Apicomplexa · Coral disease · srDNA · *Symbiodinium* · *Montastraea*

a coccidian protozoan (phylum Apicomplexa) which commonly associates with corals of the *Montastraea annularis* species complex.

Introduction

Scleractinian corals associate with diverse microbial communities (e.g., Rowan 1998; Rohwer et al. 2001). Some eukaryotic microbes are common, well-known associates of corals; notably dinoflagellates of the genus *Symbiodinium* (e.g., Trench 1987), endolithic algae (e.g., Lukas 1974), and fungi (e.g., Bentis et al. 2000). Other eukaryotic associates have not received detailed study. Descriptions of these normal associations are a necessary first step towards identifying abnormal microbial associations (i.e., pathogens) which lead to coral disease (Richardson 1998).

Methods for molecular genetic analysis have revolutionized the study of marine microbes (e.g., Brinkmeyer et al. 2000) and have proven useful in the study of coral microbial communities (e.g., Rowan 1998; Rohwer et al. 2001). Here we give a brief account of a coral-associated eukaryote which was identified by molecular methods during a bleaching experiment with the Caribbean coral *Montastraea annularis* (Toller et al. 2001a). This symbiont, provisionally named genotype N, was identified during routine screening of *Symbiodinium* samples by restriction fragment length polymorphism (RFLP) analysis of small subunit ribosomal RNA genes (srDNA). Our data indicate that genotype N represents

Methods

Genotype N was identified by an RFLP that was distinct from those of *Symbiodinium* taxa known to associate with *M. annularis* (Fig. 1a). Here, we report only those methods specific to the identification and characterization of genotype N; methods for routine RFLP analysis of *Symbiodinium* srDNA, as well as methods for experimental manipulation of corals, are given elsewhere (Toller et al. 2001a, 2001b).

Initially, the srDNA of genotype N was detected following polymerase chain reaction (PCR) with a “host-excluding” primer combination (Rowan and Powers 1991): the primers were ss5 (a universal primer) and ss3Z (a primer designed to exclude cnidarian srDNA). We first observed genotype N in samples where *Symbiodinium* cell numbers were extremely low ($<0.4 \times 10^5$ cells/cm²) following experimental bleaching of host corals (~100-fold reduction in *Symbiodinium* numbers relative to controls). These samples yielded little srDNA when PCR-amplified in the usual (Toller et al. 2001b) manner. To obtain sufficient srDNA, we used two rounds of amplification as follows: srDNA was amplified from samples with host-excluding primers for 34 PCR cycles. Aliquots (10 µl) of those amplifications were electrophoresed on agarose gels (1.0% Nuseive GTG; FMC BioProducts, Rockland, ME), and faint bands of srDNA were excised and added to 100 µl water. The gel-purified srDNA was heated to 65 °C for 2 min, and then 1 µl of each was PCR-amplified with host-excluding primers in the same manner. The resulting re-amplified srDNA was then analyzed by RFLP as described elsewhere (Toller et al. 2001b).

To further characterize genotype N, srDNA was cloned and sequenced, as described previously (Toller et al. 2001b). Samples came from three different colonies of *M. annularis* in experiment I of Toller et al. (2001a). srDNA was amplified with universal PCR primers (clone N⁰⁻¹) and with host-excluding primers (clones N⁰⁻² and N⁰⁻³). For each sample, a minimum of 12 recombinants was examined by RFLP with *Dpn* II and with *Taq* I. DNA sequences were deposited in GenBank (<http://www.ncbi.nlm.nih.gov/>) under accession numbers AF238264 (N⁰⁻¹), AF238265 (N⁰⁻²), and AF238266 (N⁰⁻³).

srDNA sequences from genotype N, *Symbiodinium* A, B, C, and E, and the coral *M. annularis* were used to design “N-specific” PCR primers that amplify only srDNA of genotype N from samples that also contained coral and zooxanthellar srDNAs: The 5′ ends of primer 18N-F2 (5′-TAGGAATCTAAACCTCTTCCA-3′; forward primer) and of primer 18N-R1 (5′-CAGGAACAAGGGTCCCCGACC-3′;

W.W. Toller (✉) · N. Knowlton
Marine Biology Research Division, 0202, Scripps Institution
of Oceanography, University of California San Diego,
La Jolla, California 92093-0202, USA
E-mail: westoller@hotmail.com
Tel.: +1-858-8222486
Fax: +1-858-5347313

R. Rowan
University of Guam Marine Laboratory,
Mangilao, Guam 96923, USA




Fig. 1. **a** RFLP analysis of srDNA from genotype N and *Symbiodinium*. srDNA was amplified with host-excluding PCR primers from a coral sample (N) and from different srDNA clones representing *Symbiodinium A* (A), *Symbiodinium B* (B), *Symbiodinium C* (C), and *Symbiodinium E* (E). srDNA was digested with *Dpn II* (left) or *Taq I* (right). The sample containing RFLP genotype N was taken from *Montastraea annularis* following experimental bleaching (see text). Lanes marked M contain size standards of (from top to bottom) 1,500 bp, 1,200 bp, and then 1,000 bp to 100 bp in 100-bp increments. **b** RFLP analysis of srDNA obtained by PCR with N-specific primers. srDNA was amplified from coral samples (lanes 1–3) and an srDNA clone (N⁰⁻¹) and digested with *Dpn II* (left) or *Taq I* (right). Samples came from three different colonies of *M. annularis*. Size markers (M) are as above

reverse primer) correspond to nucleotides 515 and 1374 (respectively) in clone N⁰⁻¹. These primers were used in PCR amplifications as before, except that DNA synthesis steps were reduced to 1 min. PCR products were analyzed by RFLP with *Dpn II* and with *Taq I*.

For phylogenetic analysis, we aligned srDNA sequences with Clustal X software (Thompson et al. 1997) and used neighbor-joining reconstruction (Saitou and Nei 1987) with 1,000 bootstrap replications. We included srDNA sequences from representative alveolate taxa (apicomplexans, dinoflagellates, ciliates) which were used previously in similar phylogenetic analyses (McNally et al. 1994; Escalante and Ayala 1995).

Results and discussion

Our initial observations of genotype N in experimentally bleached corals (see Methods) suggested that N might


normally be present at undetectable levels, and “revealed” when *Symbiodinium* numbers fall. The design of that experiment enabled us to return to archived samples taken from the same corals, but prior to bleaching (i.e., prior to reduction of *Symbiodinium* numbers). Re-analysis of these samples with N-specific PCR primers confirmed that genotype N was present (in all cases) prior to experimental treatment, when *Symbiodinium* were abundant. Using the same method, we then screened samples from other corals. Approximately 90% of samples taken from apparently healthy, unmanipulated colonies of *M. annularis* (45 colonies) and *M. faveolata* (7 colonies) tested positive for genotype N, yielding a PCR product of ca. 860 bp which, by digestion with *Dpn II* and with *Taq I*, was indistinguishable from the PCR product obtained from clone N⁰⁻¹ (Fig. 1b). These data indicate that genotype N commonly associates with *M. annularis* and *M. faveolata*, and that its presence was previously masked in our analyses by the much greater abundance of *Symbiodinium*.

Sequence data confirm that the srDNA of genotype N is quite divergent (12–17%) from those of *Symbiodinium A*, *B*, *C*, and *E* – the zooxanthellae known to associate with *M. annularis* (see Toller et al. 2001a, 2001b). Sequence comparisons also explain why genotype N was amplified with host-excluding PCR primers – the primer ss3Z has an extensive match to the srDNA of genotype N (clone N⁰⁻¹; not shown; see also McNally et al. 1994). We performed experiments with mixtures of srDNA clones from genotype N (clone N⁰⁻¹) and *Symbiodinium C* (clone C⁰; Toller et al. 2001b) and confirmed that our host-excluding protocol amplifies both srDNA with approximately equal efficiency (not shown). Thus, our method apparently detects both genotype N and *Symbiodinium* in proportion to the relative abundance of their srDNA.

Phylogenetic analysis (Fig. 2) indicates that genotype N is a member of the protozoan phylum Apicomplexa, a sister group to the dinoflagellates (Gajadhar et al. 1991; McNally et al. 1994). As far as is known, all apicomplexans are animal parasites (Levine 1982). Some, for example the apicomplexans *Plasmodium* and *Toxoplasma*, are notorious pathogens in humans and domestic animals. In our analysis, genotype N shared greatest similarity to *Toxoplasma*, *Neospora*, *Isospora*, and *Sarcocystis*, representative taxa from the apicomplexan class Coccidia (Levine 1982). From these data, we conclude that genotype N represents a coccidian protozoan that associates with corals.

Our three sequences from genotype N (from three different colonies of *M. annularis*) suggest that considerable srDNA nucleotide variation (1.0–1.7%) exists between samples obtained from different colonies. These srDNA sequences differ as much as comparisons between different species of coccidians (e.g., srDNA from *Toxoplasma gondii* and *Neospora caninum* differ by only 0.4%). However, srDNA heterogeneity is common in Apicomplexa in general (e.g., Li et al. 1997), and the accuracy and/or taxonomic significance of individual sequences may be suspect in such mixtures (see Wintz-

Fig. 2. Phylogenetic analysis of srDNA sequences from genotype N and representative protozoa. srDNA sequences were aligned and neighbor-joining reconstruction was used to explore the relationship of RFLP genotype N to members of the protozoan phyla Apicomplexa (apicomplexans, light gray box), Dinozoa (dinoflagellates, medium-gray box), and Ciliophora (ciliates, dark gray box). Bootstrap values (percent of 1,000 replicates) are shown where support for grouping exceeded 70%



ingerode et al. 1997; Toller et al. 2001b). In our limited examination of srDNA from genotype N, we found evidence of within-sample heterogeneity – approximately 25% of the srDNA clones obtained from each sample had restriction site changes not detected by RFLP of the original samples (not shown). srDNA heterogeneity within samples may contribute to the nucleotide variation that we observed between samples, and our data will serve as a starting point for clarifying this issue.

Does this coccidian associate with coral hosts outside of the *Montastraea annularis* species complex? The answer seems likely to be “yes.” Upton and Peters (1986) used light microscopy to describe a coccidium, *Gemmocystis cylindrus*, from eight species of Caribbean corals. We made a preliminary analysis of two temperate, azooxanthellate coral species (*Balanophyllia elegans* and *Astrangia* sp. collected from San Diego, CA) and they also tested positive with N-specific PCR (not shown). We suggest that the association of coccidian apicomplexans, such as genotype N, with corals may be common.

We do not yet know if genotype N is a coral parasite. Clearly, it is related to a group of totally parasitic protozoa (Fig. 2), but its ecological relationship with corals remains obscure. We detected it in corals with no outward signs of pathology. Observations on *G. cylindrus* represent an analogous case, where this presumably parasitic coccidian was only infrequently

associated with host coral pathology (Upton and Peters 1986). The use of in situ hybridization methods (e.g., Distel et al. 1995) will enable localization of N within host tissues and may further clarify the nature of their association.

Ultimately, it is likely that understanding the ecology of coral–apicomplexan associations will be more important than making a clear distinction between a benign association and a parasitic one. Coral pathogens, in some cases, may be otherwise benign microbes which become pathogenic only when their coral host becomes compromised, for example, under conditions of environmental stress (Mitchell and Chet 1975). Assays like ours can be used to rapidly assess the spatial and temporal patterns of coccidian distribution within and among host colonies. These methods can be used to test for the potential involvement of apicomplexans in coral diseases, which are of increasing concern worldwide (Richardson 1998).

Acknowledgments We thank Nick Holland, David Kline, Forest Rohwer, and three anonymous reviewers for their comments on this manuscript.

References

- Bentis CJ, Kaufman L, Golubic S (2000) Endolithic fungi in reef-building corals (Order: Scleractinia) are common, cosmopolitan, and potentially pathogenic. *Biol Bull* 198:254–260

- Brinkmeyer R, Rappé M, Gallacher S, Medlin L (2000) Development of clade- (*Roseobacter* and *Alteromonas*) and taxon-specific oligonucleotide probes to study interactions between toxic dinoflagellates and their associated bacteria. *Eur J Phycol* 35:315–329
- Distel DL, Lee HK-W, Cavanaugh CM (1995) Intracellular coexistence of methano- and thiotrophic bacteria in a hydrothermal vent mussel. *Proc Natl Acad Sci USA* 92:9598–9602
- Escalante AA, Ayala FJ (1995) Evolutionary origins of *Plasmodium* and other Apicomplexa based on rRNA genes. *Proc Natl Acad Sci USA* 92:5793–5797
- Gajadhar AA, Marquardt WC, Hall R, Gunderson J, Carmona EVA, Sogin ML (1991) Ribosomal RNA sequences of *Sarcocystis muris*, *Theileria annulata*, and *Cryptocodium cohnii* reveal evolutionary relationships among apicomplexans, dinoflagellates, and ciliates. *Mol Biochem Parasit* 45:147–154
- Levine ND (1982) Apicomplexa. In: Parker SP (ed) *Synopsis and classification of living organisms*, vol 1. McGraw-Hill, New York, pp 750–787
- Li J, Gutell RR, Damberger SH, Wirtz RA, Kissinger JC, Rogers MJ, Sattabongkot J, McCutchan TF (1997) Regulation and trafficking of three distinct 18 S ribosomal RNAs during development of the malaria parasite. *J Mol Biol* 269:203–213
- Lukas K (1974) Two species of the chlorophyte genus *Ostreobium* from skeletons of Atlantic and Caribbean reef corals. *J Phycol* 10:331–335
- McNally KL, Govind NS, Thomé PE, Trench RK (1994) Small-subunit ribosomal DNA sequence analyses and a reconstruction of the inferred phylogeny among symbiotic dinoflagellates (Pyrrophyta). *J Phycol* 30:316–329
- Mitchell R, Chet I (1975) Bacterial attack of corals in polluted seawater. *Microbial Ecol* 2:227–233
- Richardson LL (1998) Coral diseases: what is really known? *Trends Ecol Evol* 13:438–443
- Rohwer F, Breitbart M, Jara J, Azam F, Knowlton N (2001) Diversity of bacteria associated with the Caribbean coral *Montastraea franksi*. *Coral Reefs* 20:85–95
- Rowan R (1998) Diversity and ecology of zooxanthellae on coral reefs. *J Phycol* 34:407–417
- Rowan R, Powers DA (1991) Molecular genetic identification of symbiotic dinoflagellates (zooxanthellae). *Mar Ecol Prog Ser* 71:65–73
- Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4:406–425
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997) The Clustal X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* 24:4876–4882
- Toller WW, Rowan R, Knowlton N (2001a) Repopulation of zooxanthellae in the Caribbean corals *Montastraea annularis* and *M. faveolata* following experimental and disease-associated bleaching. *Biol Bull* 201:360–373
- Toller WW, Rowan R, Knowlton N (2001b) Zooxanthellae of the *Montastraea annularis* species complex: patterns of distribution of four taxa of *Symbiodinium* on different reefs and across depths. *Biol Bull* 201:348–359
- Trench R (1987) Dinoflagellates in non-parasitic symbioses. In: Taylor FJR (ed) *The biology of dinoflagellates*. Botanical Monographs vol 21. Blackwell, Oxford, pp 531–570
- Upton SJ, Peters EC (1986) A new and unusual species of coccidium (Apicomplexa: Agamococcidiorida) from Caribbean scleractinian corals. *J Invert Pathol* 47:184–193
- Wintzingerode Fv, Göbel UB, Stackebrandt E (1997) Determination of microbial diversity in environmental samples: pitfalls of PCR-based rRNA analysis. *FEMS Microbiol Rev* 21:213–229